

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/14, 487/04, 513/04, 471/04, 495/04, 401/04, A61K 31/445		A1	(11) International Publication Number: WO 96/23784
			(43) International Publication Date: 8 August 1996 (08.08.96)
(21) International Application Number: PCT/EP96/00363		(81) Designated States: AL, AM, AU, BB, BG, BR, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KR, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KG, KZ, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 23 January 1996 (23.01.96)			
(30) Priority Data: 95200229.3 31 January 1995 (31.01.95) EP (34) Countries for which the regional or international application was filed: DE et al.			
(71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE).		<p>Published</p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
(72) Inventors; and (75) Inventors/Applicants (for US only): VANDENBERK, Jan [BE/BE]; Kempenlaan 15, B-2340 Beerse (BE). KENNIS, Ludo, Edmond, Josephine [BE/BE]; Guido Gezellestraat 50, B-2300 Turnhout (BE). MERTENS, Josephus, Carolus [BE/BE]; Staatsbaan 35/1, B-2360 Oud-Turnhout (BE).			
(74) Common Representative: JANSSEN PHARMACEUTICA N.V.; Patent Dept., Turnhoutseweg 30, B-2340 Beerse (BE).			
(54) Title: ANTIPSYCHOTIC 4-(1H-INDOLYL-1-YL)-1-SUBSTITUTED PIPERIDINE DERIVATIVES			
<div style="text-align: center;"> <p>(I)</p> </div>			
(57) Abstract			
<p>The present invention concerns the compounds of formula (I), the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein the dashed line designates an optional bond; R¹ and R² are each independently hydrogen, halogen, C₁-alkyl or C₁-alkyloxy; R³ and R⁴ are each independently hydrogen, C₁-alkyl, phenyl or phenyl substituted with one, two or three substituents selected from halo, hydroxy, nitro, cyano, trifluoromethyl, C₁-alkyl, C₁-alkyloxy, C₁-alkylthio, mercapto, amino, mono- and di(C₁-alkyl)-amino, carboxyl, C₁-alkyloxycarbonyl and C₁-alkylcarbonyl; Alk is C₁-alkanediyl; D is a pyrimidinone, piperidone or a benzimidazolidinone; having antipsychotic activity; their preparation, compositions containing them and their use as a medicine.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

ANTIPSYCHOTIC 4-(1H-INDOLYL-1-YL)-1-SUBSTITUTED PIPERIDINE DERIVATIVES

The present invention concerns 4-(1H-indol-1-yl)-1-piperidiny derivatives having therapeutic potential in psychotic disorders. It further relates to their preparation, compositions comprising them and their use as a medicine.


In EP-A-0,037,265, published on October 7, 1981, EP-A-0,070,053, published on January 19, 1983, and in EP-A-0,378,255, published on July 18, 1990, there are described a number of 5-[[4-(1H-indol-3-yl)-1-piperidinyl]alkyl]-4(3H)-pyrimidinone derivatives having antipsychotic, antihistaminic and antiserotonergic activity.

Structurally, the compounds of the present invention differ therefrom in that the 4 position of their piperidine moiety is invariably substituted by a 1H-indol-1-yl derivative.

EP-A-0,470,039, published on February 5, 1992, discloses 4-(3-aryl-1H-indol-1-yl)-1-piperidiny derivatives as selective antagonists of the serotonin 5-HT₂ receptor without substantial dopamine D-2 antagonistic activity both *in vivo* and *in vitro*. Said selective antagonistic property is measured as the ratio between the dopamine D-2 receptor and the serotonin 5-HT₂ receptor antagonistic activities. Unexpectedly, the compounds of the present invention differ therefrom in that they exhibit central dopamine and central serotonin antagonistic activity *in vivo* in similar dose-ranges, i.e. the ratio between central dopaminergic and central serotonergic activity is about unity.

The compounds of the present invention show antipsychotic activity with an unexpected increased cardiovascular safety, i.e. they show an improved dissociation between the peripheral α -adrenergic antagonistic activity and the central dopamine and serotonin antagonistic activity.

The present invention concerns the compounds of formula



the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein :

the dashed line designates an optional bond;

R¹ and R² are each independently hydrogen, halogen, C₁-alkyl or C₁-alkyloxy;


-2-

R^3 and R^4 are each independently hydrogen, C₁-6alkyl, phenyl or phenyl substituted with one, two or three substituents selected from halo, hydroxy, nitro, cyano, trifluoromethyl, C₁-6alkyl, C₁-6alkyloxy, C₁-6alkylthio, mercapto, amino, mono-

and di(C₁-6alkyl)amino, carboxyl, C₁-6alkyloxycarbonyl and C₁-6alkylcarbonyl;

Alk is C₁-4alkanediyl;

D is a radical of formula



wherein R^5 is hydrogen or C₁-6alkyl;

R^6 is hydrogen, C₁-6alkyl, C₁-6alkyloxy, C₁-6alkylthio, amino or mono- or di(C₁-6alkyl)amino; and

R^7 is hydrogen or C₁-6alkyl; or

R^6 and R^7 taken together may form a bivalent radical $-R^6-R^7-$, in particular, $-R^6-R^7-$ may be

$-CH_2-CH_2-CH_2-$ (a-1);

$-CH_2-CH_2-CH_2-CH_2-$ (a-2);

$-CH=CH-CH_2-$ (a-3);

$-CH_2-CH=CH-$ (a-4) or

$-CH=CH-CH=CH-$ (a-5);

wherein one or two hydrogen atoms of said radicals (a-1) to (a-5) each independently may be replaced by C₁-6alkyl, hydroxy, C₁-6alkyloxy or C₁-10alkylcarbonyloxy; or

$-R^6-R^7-$ may also be

$-S-CH_2-CH_2-$ (a-6);

$-S-CH_2-CH_2-CH_2-$ (a-7);

$-S-CH=CH-$ (a-8);

$-NH-CH_2-CH_2-$ (a-9);

$-NH-CH_2-CH_2-CH_2-$ (a-10);

$-NH-CH=CH-$ (a-11);

$-NH-CH=N-$ (a-12);


$-S-CH=N-$ (a-13) or

$-CH=CH-O-$ (a-14);

wherein one or where possible two or three hydrogen atoms in said radicals (a-6) to (a-14) each independently may be replaced by C₁-6alkyl;

or D is a radical of formula

-3-



wherein R^8 is hydrogen or C_{1-6} alkyl.

As used in the foregoing definitions the term halogen is generic to fluoro, chloro, bromo and iodo. The term C_{1-4} alkyl defines straight and branched saturated hydrocarbons, having from 1 to 4 carbon atoms such as, for example, methyl, ethyl, propyl, butyl, 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl. The term C_{1-6} alkyl is meant to include C_{1-4} alkyl radicals and the higher homologues thereof having 5 or 6 carbon atoms such as, for example, pentyl, hexyl and the like. The term C_{1-10} alkyl is meant to include C_{1-6} alkyl radicals and the higher homologues thereof having 7 to 10 carbon atoms such as, for example, heptyl, octyl, nonyl, decyl and the like. The term C_{2-4} alkanediyl defines bivalent straight or branch chained alkanediyl radicals having from 2 to 4 carbon atoms such as, for example, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl and the like. The term C_{1-4} alkanediyl is meant to include C_{2-4} alkanediyl radicals and the lower homologue, i.e. 1,1-methanediyl.

The addition salts as mentioned herein are meant to comprise the therapeutically active addition salt forms which the compounds of formula (I) are able to form with appropriate acids, such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

The pharmaceutically acceptable addition salts as mentioned hereinabove are also meant to comprise the therapeutically active non-toxic base, in particular, a metal or amine addition salt forms which the compounds of formula (I) are able to form. Said salts can conveniently be obtained by treating the compounds of formula (I) containing acidic hydrogen atoms with appropriate organic and inorganic bases such as, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-*D*-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

Conversely said salt forms can be converted by treatment with an appropriate base or acid into the free acid or base form.


- 5 The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) are able to form and said solvates are meant to be included within the scope of the present invention. Examples of such solvates are, e.g. the hydrates, alcoholates and the like.

- 10 The term stereochemically isomeric forms as used herein defines all the possible isomeric forms in which the compounds of formula (I) may occur. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure.


- 15 Some of the compounds of formula (I) may also exist in their tautomeric forms. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

- 20 Whenever used hereinafter, the term compounds of formula (I) is meant to include also the pharmaceutically acceptable addition salts and all stereoisomeric forms.

Interesting compounds are those compounds of formula (I) wherein D is a radical of formula



-5-



(d-10)

wherein R^5 is hydrogen or C₁-6alkyl and preferably is methyl; R^9 and R^{10} each independently are hydrogen or C₁-6alkyl; R^{11} is C₁-6alkyl, hydroxy, C₁-6alkyloxy or C₁-10alkylcarbonyloxy, and preferably is hydroxy, C₁-6alkyloxy or C₁-10alkylcarbonyloxy; and R^{12} is hydrogen, C₁-6alkyl, C₁-6alkyloxy, C₁-6alkylthio, amino or mono- or di(C₁-6alkyl)amino.

Further interesting compounds are those compounds of formula (I) wherein R^3 and R^4 are both hydrogen.

In a first subset of compounds of formula (I), Alk is C₂-4alkanediyl.

In a second subset of compounds of formula (I), R^1 is hydrogen or halogen and R^2 is hydrogen.

In a third subset of compounds of formula (I), the substituent on the 4 position of the piperidiny is an 1*H*-indole, i.e. the dashed line represents an extra bond.


In a fourth subset of compounds of formula (I), the substituent on the 4 position of the piperidiny is an 2,3-dihydro-1*H*-indole.

Preferred compounds are those compounds of formula (I) wherein R^1 is hydrogen or fluor; R^2 , R^3 and R^4 are hydrogen; Alk is 1,2-ethanediyl; D is a radical of formula (a) wherein R^5 is methyl; R^6 is C₁-6alkylamino and R^7 is C₁-6alkyl; or R^6 and R^7 taken together may form a bivalent radical of formula (a-2) or (a-5) wherein one hydrogen atom of said radicals (a-2) and (a-5) may be replaced by C₁-6alkyl, hydroxy or C₁-6alkyloxy; or R^6 and R^7 taken together may form a bivalent radical of formula (a-6), (a-7), (a-8), (a-13) or (a-14) wherein one or where possible two hydrogen atoms may be replaced by C₁-6alkyl.

More preferred compounds are those preferred compounds wherein the substituent on the 4 position of the piperidiny is an indole wherein R^1 is substituted in the 5 position and stands for hydrogen or fluor.

- Most preferred compounds are selected from
- 3-[2-[4-(5-fluoro-1*H*-indol-1-yl)-1-piperidinyl]ethyl]-6,7,8,9-tetrahydro-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;
- 6-[2-[4-(5-fluoro-1*H*-indol-1-yl)-1-piperidinyl]ethyl]-7-methyl-5*H*-thiazolo[3,2-*a*]-pyrimidin-5-one;
- 5 5-[2-[4-(5-fluoro-1*H*-indol-1-yl)-1-piperidinyl]ethyl]-3,6-dimethyl-2-(methylamino)-4(3*H*)-pyrimidinone; and
- 2,3-dihydro-6-[2-[4-(1*H*-indol-1-yl)-1-piperidinyl]ethyl]-7-methyl-5*H*-thiazolo[3,2-*a*]-pyrimidin-5-one;
- 10 the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof.


- The compounds of formula (I) can generally be prepared by *N*-alkylating a 1-(piperidin-4-yl)-1*H*-indole derivative of formula (II) with an alkylating reagent of formula (III) following the procedure described in EP-A-0,037,265, EP-A-0,070,053,
- 15 EP-A-0,196,132 and in EP-A-0,378,255.



- 20 In the following intermediates, the dashed line and the radicals D, Alk and R¹ to R⁴ are defined as under formula (I) unless otherwise specified. In intermediate (III), W represents an appropriate reactive leaving group such as, for example, halo, e.g. chloro, bromo or iodo; sulfonyloxy, e.g. methanesulfonyloxy, 4-methylbenzenesulfonyloxy.
- 25 In this and the following preparations, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, trituration and chromatography.
- 30 Compounds of formula (I) wherein R³ and R⁴ are hydrogen and the heterocyclic moiety in the 4 position of the piperidine ring is an 1*H*-indole, said compounds being represented by formula (I-f), may alternatively be prepared by cyclizing an intermediate of formula (IV) wherein W is a reactive leaving group such as, for example, halo, e.g. chloro, bromo or iodo; sulfonyloxy, e.g. methanesulfonyloxy, 4-methylbenzene-

-7-

sulfonyloxy, in the presence of a reductant such as, for example, a complex metal hydride, e.g. sodium borohydride. Said procedure is described in Synthetic Communications, 18(3), 265-273 (1988).



EP-A-0,037,265, EP-A-0,070,053, EP-A-0,196,132 and EP-A-0,378,255 describe other alternative procedures to prepare compounds of formula (I). Furthermore, the compounds of formula (I) may be converted into each other following art-known functional group transformation reactions.


10

A number of intermediates and starting materials are known compounds which may be prepared according to art-known methodologies. For example, intermediates of formula (III) and their preparations are described in EP-A-0,037,265, EP-A-0,070,053, EP-A-0,196,132 and in EP-A-0,378,255.

15

Intermediates of formula (II) can be prepared following art-known procedures, e.g. by reductively aminating a ketone of formula (V), wherein P is a protecting group such as, for example, an alkoxycarbonyl group, with an indole derivative of formula (VI), and subsequently removing the protecting group P by art-known deprotection techniques.

20



Said reductive amination may conveniently be carried out by mixing the reactants in a suitable reaction-inert solvent such as, for example, methanol or toluene, with an appropriate reductant. Preferably, the ketone of formula (V) is first reacted with the indole derivative of formula (VI) to form an enamine, which is subsequently reduced. Hydrogen in the presence of a suitable catalyst such as, for example, palladium or platinum supported on for instance charcoal may be used as an appropriate reductant.

25

Elevated pressure and/or temperature may enhance the rate of the reaction. In order to prevent the undesired further hydrogenation of certain functional groups in the reactants and the reaction products, it may be advantageous to add an appropriate catalyst poison to the reaction mixture such as, for example, thiophene.

The intermediates of formula (VI) wherein the dashed line represents an extra bond and R^3 and R^4 are both hydrogen, may be prepared as described in Journal of Heterocyclic Chemistry, 2, 298-299 (1965).

Some of the compounds of formula (I) and some of the intermediates in the present invention contain at least one asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or compounds by, for example, selective crystallization or chromatographic techniques, e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers.

Pure stereochemically isomeric forms of the compounds of formula (I) may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically. The pure and mixed stereochemically isomeric forms of the compounds of formula (I) are intended to be embraced within the scope of the present invention.

The compounds of formula (I), the pharmaceutically acceptable addition salts and stereochemically isomeric forms thereof, are antagonists of neurotransmitters and in particular of the mediators serotonin and dopamine. Antagonizing said mediators will suppress or relieve a variety of symptoms associated with phenomena induced by the release, in particular the excessive release, of these mediators. Therapeutic indications for using the present compounds are mainly in the CNS area, especially in psychotic disorders. Further, serotonin is a potent broncho- and vasoconstrictor and thus the present antagonists may be used against hypertension and vascular disorders. In addition, serotonin antagonists have been associated with a number of other properties such as, the suppression of appetite and promotion of weight loss, which may prove

effective in combatting obesity; and also the alleviation of withdrawal symptoms in addicts trying to discontinue drinking and smoking habits. The present compounds also appear to be useful therapeutic agents for combatting autism.

- 5 In view of the usefulness of the subject compounds in the treatment of diseases associated with the release of neurotransmitters, in particular in the treatment of psychotic diseases, the present invention provides a method of treating warm-blooded animals suffering from such diseases, in particular psychotic diseases, said method comprising the systemic administration of an antipsychotic amount of a compound of formula (I) or a
10 pharmaceutically acceptable addition salt thereof, effective in treating diseases associated with the release of neurotransmitters, in particular psychotic diseases. Those of skill in the treatment of such diseases could readily determine the effective amount from the test results presented hereinafter. In general it is contemplated that an effective antipsychotic daily amount would be from about 0.01 mg/kg to about 4 mg/kg body weight, more
15 preferably from about 0.04 mg/kg to about 2 mg/kg body weight.

The present invention thus also relates to compounds of formula (I) as defined hereinabove for use as a medicine. Further, the present invention also relates to the use of a compound of formula (I) for the manufacture of a medicament for treating psychotic
20 diseases.

- The "apomorphine, tryptamine, norepinephrine (ATN) test in rats" is used to evaluate dopamine antagonism, serotonin antagonism and α -adrenergic antagonistic properties of the compounds of formula (I). In said test, which is described hereinafter, rats are
25 observed for effects which are indicative for peripheral and central activity of the tested compounds. Centrally acting serotonin antagonists are potential antipsychotic drugs, in particular when simultaneously displaying dopamine antagonism. Peripheral serotonin antagonists are potentially useful in the gastro-intestinal and cardiovascular field, in particular when simultaneously displaying α -adrenergic antagonistic activity. The
30 compounds of the present invention show a strong central dopamine and serotonin antagonism and little peripheral α -adrenergic antagonistic activity. In view of the central dopamine and serotonin antagonistic activity of the compounds of formula (I), they are particularly useful in combatting psychoses, aggressive behaviour, anxiety, depression and migraine. Furthermore, the improved dissociation between peripheral α -adrenergic
35 antagonistic activity and central dopamine and serotonin antagonistic activity of the compounds of formula (I) over prior known antipsychotic agents, leads to an increased cardiovascular safety, i.e. a decreased likelihood of hypotension.

For administration purposes, the subject compounds may be formulated into various pharmaceutical forms. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, in addition salt or in free acid or base form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable solutions containing compounds of formula (I) may be formulated in an oil for prolonged action. Appropriate oils for this purpose are, for example, peanut oil, sesame oil, cottonseed oil, corn oil, soy bean oil, synthetic glycerol esters of long chain fatty acids and mixtures of these and other oils. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wettable agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause any significant deleterious effects on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on or as an ointment. Addition salts of (I) due to their increased water solubility over the corresponding free base or free acid form, are obviously more suitable in the preparation of aqueous compositions.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active

ingredient calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

The following examples are intended to illustrate and not to limit the scope of the present invention.

10 Experimental part

Hereinafter, "RT" means room temperature and "DIPE" means diisopropyl ether.

A. Preparation of the intermediates

Example 1

- 15 a) Sodium methylate (1260.5 g) was added dropwise to a suspension of thiourea (373 g) in ethanol (1750 ml) and the dropping funnel was rinsed with ethanol (175 ml). The mixture was heated to 40 °C and 3-acetyldihydro-2(3H)-furanone (448.5 g) was added dropwise. The dropping funnel was rinsed with ethanol (475 ml) and the reaction mixture was stirred and refluxed for 6.25 hours. Stirring was continued for 2 hours,
- 20 while cooling. The solvent was evaporated and the residue was dissolved in water (2800 ml). This mixture was cooled on ice and neutralized with hydrochloric acid (490 ml). The precipitate was filtered off, washed with water (350 ml) and dried, yielding 268 g (41.1%) of 5-(2-hydroxyethyl)-2-mercapto-6-methyl-4(3H)-pyrimidinone (intermediate-1).
- 25 b) A mixture of intermediate 1 (37.2 g) and sodium methylate (36 g) in methanol (250 ml) was stirred for 30 min. Ethane iodide (31.2 g) was added dropwise. The reaction mixture was stirred and refluxed for 3 hours. The solvent was evaporated and the residue was stirred in water, filtered and recrystallized from CH₃CN. The precipitate was filtered off and dried, yielding 26 g (60%) of 2-(ethylthio)-5-(2-hydroxyethyl)-6-
- 30 methyl-4(3H)-pyrimidinone (intermediate 2).
- c) Potassium hydroxide (8.4 g) was added to a solution of intermediate 2 (33 g) in dimethyl sulfoxide (150 ml). The mixture was stirred for 1 hour at 60-70 °C. The mixture was cooled to 20 °C and iodomethane (21.3 g) was added dropwise. The reaction mixture was stirred overnight at RT. The mixture was poured out into water and
- 35 extracted with toluene. The separated organic layer was dried, filtered and the solvent was evaporated. The residue was crystallized from 4-methyl-2-pentanone and the precipitate was filtered off and dried, yielding 16 g (46%) of 2-(ethylthio)-5-(2-hydroxyethyl)-3,6-dimethyl-4(3H)-pyrimidinone (intermediate 3).

Example 2

- a) 3-acetyldihydro-2(3*H*)-furanone (25.6 g) was added to a stirred mixture of acetamidine hydrochloride (20 g) and sodium methylate (110 g) in methanol (150 ml). The reaction mixture was stirred and refluxed overnight. The reaction mixture was cooled and acetic acid (36 g) was added dropwise. The precipitate was filtered off and the filtrate evaporated. The residue was purified by column chromatography over silica gel (eluent: CHCl₃/(CH₃OH/NH₃) 90/10). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from 2-propanol. The crystals were filtered off and dried, yielding 7.6 g (22%) of 5-(2-hydroxyethyl)-2,6-dimethyl-4(3*H*)-pyrimidinone (intermediate 4).
- b) Sodium methylate (1.9 g) was added to a mixture of intermediate 4 (6 g) in methanol (50 ml) while stirring at RT. Stirring was continued for 30 minutes and iodomethane (5 g) was added dropwise and the reaction mixture was stirred and refluxed for 4 hours. The solvent was evaporated, water was added to the residue and this mixture was extracted with CH₂Cl₂. The organic layer was separated, dried over MgSO₄, filtered and the solvent was evaporated, yielding 6.0 g (94%) of 5-(2-hydroxyethyl)-2,3,6-trimethyl-4(3*H*)-pyrimidinone (intermediate 5).

Example 3

- a) A mixture of ethyl 4-oxo-1-piperidinecarboxylate (85 g) and 2,3-dihydro-1*H*-indole (60 g), palladium on activated carbon (10 %) (4 g) and a solution of thiophene in isopropyl ether (4 %) (2 ml) in methanol (700 ml) was reacted in a Parr Pressure Vessel at 50 °C overnight. After completion, the mixture was filtered and the filtrate was evaporated, yielding 80 g (58 %) of ethyl 4-(2,3-dihydro-1*H*indol-1-yl)-1-piperidinecarboxylate (intermediate 6).
- b) A mixture of intermediate 6 (95 g) and potassium hydroxide (194 g) in 2-propanol (1300 ml) was stirred and refluxed for 24 hours. The solvent was evaporated and the residue was dissolved in H₂O/CH₂Cl₂. The organic layer was separated, dried over MgSO₄, filtered and evaporated, yielding 64 g (90.4%) of 2,3-dihydro-1-(4-piperidiny)-1*H*-indole. A sample (4.5 g) was crystallized from CH₃OH and converted into the hydrochloric acid salt (1:1) in 2-propanol and filtered off, yielding 3.0 g of 2,3-dihydro-1-(4-piperidiny)-1*H*-indole monohydrochloride (intermediate 7).

Example 4

- A mixture of 2-bromoethanol (3.75 g), 1-(4-piperidiny)-1*H*-indole (5 g) and sodium hydrogen carbonate (4.2 g) in ethanol (100 ml) was stirred and refluxed overnight. The solvent was evaporated. The residue was stirred in water and this mixture was extracted with CH₂Cl₂. The separated organic layer was dried over MgSO₄, filtered and the

solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was stirred in DIPE and the solvent was evaporated, yielding 4 g (64%) of 4-(1*H*-indol-1-yl)-1-piperidineethanol (intermediate 8).

Example 5

a) A mixture of 2-amino-3-pyridinol (100 g), 3-acetyldihydro-2(3*H*)-furanone (100 g), 4-methylbenzene sulfonic acid (1 g) and xylene (700 ml) was stirred and refluxed overnight using a water separator. The mixture was cooled and the product was filtered off and dried. The product was converted into the hydrochloric acid salt in 2-propanol. The salt was filtered off and dried, yielding 120 g (58.4%) of 9-hydroxy-3-(2-hydroxyethyl)-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one monohydrochloride (intermediate 9).

b) Dimethyl sulfate (2.52 g) was added dropwise to a mixture of intermediate 9 (4.4 g) and sodium hydroxide (0.8 g) in water (10 ml), while cooling in ice water. The reaction mixture was stirred for 15 minutes at RT, then it was heated for 1 hour using a warm water bath. The reaction mixture was cooled and extracted with CH₂Cl₂. The precipitate in the separated aqueous layer was filtered off and purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 97/3). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from CH₃CN.

The crystals were filtered off and dried, yielding 2 g (42 %) of 3-(2-hydroxyethyl)-9-methoxy-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (intermediate 10).

c) A mixture of intermediate 10 (14 g) in methanol (250 ml) was hydrogenated at 50 °C with palladium on activated carbon (10 %) (2 g) as a catalyst. After uptake of hydrogen (2 eq.), the catalyst was filtered off. The filtrate was evaporated and the residue was purified by column chromatography over silica gel (eluent : CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated, yielding 10 g (71 %) of (±)-6,7,8,9-tetrahydro-3-(2-hydroxyethyl)-9-methoxy-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (intermediate 11).

Example 6

Methanesulfonyl chloride (3.43 g) was added dropwise to a stirred and cooled (ice water bath) mixture of intermediate 13 (7 g) and *N,N*-diethylethanamine (3 g) in dichloromethane (50 ml). The reaction mixture was stirred for 2 hours at RT. The reaction mixture was washed with water and the organic layer was separated, dried over MgSO₄, filtered and the solvent was evaporated, yielding 8 g (84%) of (±)-6,7,8,9-tetrahydro-9-methoxy-2-methyl-3-[2-[(methylsulfonyl)oxy]ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (intermediate 12).

In a similar way, 2-(ethylthio)-3,6-dimethyl-5-[2-(methylsulfonyl)oxy]ethyl]-4(3*H*)-pyrimidinone hemihydrate (intermediate 13) was prepared from intermediate 3 and 2,3,6-trimethyl-5-[2-(methylsulfonyl)oxy]ethyl]-4(3*H*)-pyrimidinone (intermediate 14) was prepared from intermediate 5.

Example 7

a) A mixture of intermediate 9 (170 g) and hydrobromic acid (48 %) (1000 ml) was stirred and refluxed overnight. The mixture was cooled and the precipitate was filtered off, yielding a first fraction of 100 g. The filtrate was evaporated, yielding a second fraction of 30 g. Both fractions were recrystallized from water, yielding 75 g (34 %) of 3-(2-bromoethyl)-9-hydroxy-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one monohydrobromide (intermediate 15).

b) A mixture of intermediate 15 (55 g) and hydrobromic acid (48 %) (55 g) in methanol (700 ml) was hydrogenated with palladium on activated carbon (10 %) (5 g) as a catalyst. After completion, the precipitate was filtered off and the filtrate evaporated. Ammonium hydroxide was added to the residue which was subsequently extracted with CHCl_3 . The separated organic layer was dried and evaporated. The residue was purified by column chromatography over silica gel (eluent : $\text{CHCl}_3/\text{CH}_3\text{OH}$ 90/10). The pure fractions were collected, the solvent was evaporated and the residue was recrystallized from 4-methyl-2-pentanone, yielding 17 g (44 %) of (\pm)-6,7,8,9-tetrahydro-3-(2-bromoethyl)-9-hydroxy-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (intermediate 16).

B. Preparation of the compounds of formula (I)

Example 8

A mixture of 1-(4-piperidiny)-1*H*-indole (4.1 g), 6-(2-chloroethyl)-7-methyl-5*H*-thiazolo[3,2-*a*]pyrimidin-5-one (3.4 g), prepared as described in EP-A-0,196,132, sodium carbonate (6 g) and potassium iodide (0.1 g) in 4-methyl-2-pentanone (250 ml) was stirred and refluxed overnight. The warm reaction mixture was filtered and the filtrate was evaporated. The residue was purified over silica gel on a glass filter (eluent : $\text{CH}_2\text{Cl}_2/\text{C}_2\text{H}_5\text{OH}$ 90/10). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from DIPE with a small amount of CH_3CN . The precipitate was filtered off and dried, yielding 3 g (51%) 6-[2-[4-(1*H*-indol-1-yl)-1-piperidiny]ethyl]-7-methyl-5*H*-thiazolo[3,2-*a*]pyrimidin-5-one (compound 3; mp. 134.4 °C).

Example 9

6-[2-[4-(1*H*-indol-1-yl)-1-piperidinyl]ethyl]-3,7-dimethyl-5*H*-thiazolo[3,2-*a*]pyrimidin-5-one was prepared according to the procedure described in example 7 without the presence of potassium iodide (compound 6; mp. 189.8 °C).

Example 10

A mixture of 6-(2-chloroethyl)-2,7-dimethyl-5*H*-1,3,4-thiadiazolo[3,2-*a*]pyrimidin-5-one (3.6 g), prepared as described in EP-A-0,353,821, 5-fluoro-1-(4-piperidinyl)-1*H*-indole (3.3 g), prepared according to the procedure described in Synthetic Communications, 18 (3), 265-273 (1988), sodium carbonate (1 g), potassium iodide (catalytic quantity) and sodium hydrogen carbonate (2 g) in 4-methyl-2-pentanone (150 ml) and 2-ethoxyethanol (50 ml) was stirred and refluxed overnight. The mixture was filtered and the filtrate was washed with water. The organic layer was separated, dried, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from DIPE/CH₃CN. The precipitate was filtered off, dried and purified by high-performance liquid chromatography over RP-18 (eluent: (0.5% NH₄OAc in H₂O)/CH₃OH/THF 50/40/10). The pure fractions were collected and the solvent was evaporated, yielding 1.37 g (20%) 6-[2-[4-(5-fluoro-1*H*-indol-1-yl)-1-piperidinyl]-ethyl]-2,7-dimethyl-5*H*-1,3,4-thiadiazolo[3,2-*a*]pyrimidin-5-one (compound 8; mp. 194.1 °C).

Example 11

A mixture of 6-(2-bromoethyl)-1,2,7-trimethyl-1*H*,5*H*-imidazo[1,2-*a*]pyrimidin-5-one (8.85 g), prepared as described in EP-0,378,255, 1-(4-piperidinyl)-1*H*-indole (2 g) and *N*-(1-methylethyl)-2-propanamine (1.2 g) in ethanol (100 ml) was stirred and refluxed for 6 hours. The solvent was evaporated and the residue was stirred in water and subsequently extracted with CH₂Cl₂. The separated organic layer was dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from CH₃CN/DIPE and the precipitate was filtered off and dried, yielding 2.5 g (62%) 6-[2-[4-(1*H*-indol-1-yl)-1-piperidinyl]ethyl]-1,2,7-trimethylimidazo[1,2-*a*]pyrimidin-5(1*H*)-one (compound 9; mp. 165.7 °C).

Example 12

1-(2-bromoethyl)-2*H*-benzimidazol-2-one (4.8 g) was added to a mixture of 1-(4-piperidinyl)-1*H*-indole (5.6 g), sodium methylate (1 g) and sodium carbonate (5 g) in 4-methyl-2-pentanone (250 ml). The reaction mixture was stirred and refluxed

overnight. The mixture was filtered while still warm. The filtrate was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH₂Cl₂/C₂H₅OH 90/10).


- The pure fractions were collected and the solvent was evaporated. The residue was dissolved in warm ethanol and converted into the (E)-2-butenedioic acid salt (1:1). The mixture was cooled. The precipitate was filtered off and dried, yielding: 8.7 g (91%) 1,3-dihydro-1-[2-[4-(1*H*-indol-1-yl)-1-piperidinyl]ethyl]-2*H*-benzimidazol-2-one (E)-2-butenedioate (1:1) (compound 26; mp. 205.3 °C)

Example 13

- A mixture of 5-fluoro-1-(4-piperidinyl)-1*H*-indole (3.2 g), 1-(3-chloropropyl)-1,3-dihydro-2*H*-benzimidazol-2-one (3 g), *N,N*-diethylethanamine (3 ml) and potassium iodide (0.1 g) in toluene (200 ml) was stirred and refluxed overnight. The reaction mixture was cooled, washed with water and the organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was dissolved in ethanol and converted into the (E)-2-butenedioic acid salt (1:1). The mixture was allowed to cool, while stirring. The precipitate was filtered off and dried, yielding 3 g (42%) 1-[3-[4-(5-fluoro-1*H*-indol-1-yl)-1-piperidinyl]propyl]-1,3-dihydro-2*H*-benzimidazol-2-one (E)-2-butenedioate (1:1) (compound 28; mp. 192.5 °C).

Tables 1 to 4 list similar compounds that were prepared according to one of the above described examples.

Table 1




Co. No.	Ex. No.	R ¹	R ³	R ⁴	-R ⁶ -R ⁷ -	Phys. Data (mp in °C)
1	9	H	H	H	-S-CH ₂ -CH ₂ -	190.2
2	9	F	H	H	-S-CH ₂ -CH ₂ -	200.0
3	8	H	H	H	-S-CH=CH-	134.4
4	8	F	H	H	-S-CH=CH-	126.6
6	9	H	H	H	-S-CH=C(CH ₃)-	189.8

-17-


Co. No.	Ex. No.	R ¹	R ³	R ⁴	-R ⁶ -R ⁷ -	Phys. Data (mp in °C)
8	10	F	H	H	-S-C(CH ₃)=N-	194.1
9	11	H	H	H	-N(CH ₃)-C(CH ₃)=CH-	165.7
10	8	F	H	H	-CH=C(CH ₃)-O-	208.4
11	8	H	H	H	-(CH ₂) ₄ -	142.3
12	8	F	H	H	-(CH ₂) ₄ -	173.8
13	11	H	H	H	-CH(OH)-(CH ₂) ₃ -	(±) / 154.4
14	11	F	H	H	-CH(OH)-(CH ₂) ₃ -	(±) / 192.0
15	11	F	H	H	-CH(OCH ₃)-(CH ₂) ₃ -	(±) / 159.6
16	9	H	H	H	-S-(CH ₂) ₃ -	153.0
17	9	F	H	H	-S-(CH ₂) ₃ -	173.1
18	13	F	H	H	-CH=CH-CH=CH-	144.4
19	8	H	H	H	-C(CH ₃)=CH-CH=CH-	127.6
20	8	F	H	H	-C(CH ₃)=CH-CH=CH-	165.7
29	11	F	H	H	-(CH ₂) ₃ -CH(CH ₃)-	165.9
30	11	F	-CH ₃	-C ₆ H ₅	-(CH ₂) ₄ -	147.7
32	11	H	-CH ₃	H	-(CH ₂) ₄ -	140.2
33	11	-OCH ₃	H	H	-(CH ₂) ₄ -	154.7
36	11	H	-CH ₃	-CH ₃	-(CH ₂) ₄ -	149.2
37	11	-CH ₃	H	H	-(CH ₂) ₄ -	161.2
38	11	Cl	H	H	-(CH ₂) ₄ -	184.0

Table 2




Co. No.	Ex. No.	R ¹	R ³	R ⁴	-R ⁶ -R ⁷ -	Phys. Data (mp in °C)
5	11	H	H	H	-S-CH=CH-	143.4
31	11	H	H	H	-(CH ₂) ₄ -	119.2
34	11	H	-CH ₃	H	-(CH ₂) ₄ -	99.4
35	11	H	-CH ₃	-CH ₃	-(CH ₂) ₄ -	123.7

-18-

Table 3

Co. No.	Ex. No.	R ¹	R ⁶	Physic. Data (mp. in °C)
21	11	H	-CH ₃	119.5
22	11	H	-NH ₂	210.7
23	8	H	-NH-CH ₃	171.9
24	8	F	-NH-CH ₃	179.8
25	11	H	-S-CH ₂ -CH ₃	129.5

5 Table 4

Co. No.	Ex. No.	R ¹	n	x	Phys. Data (mp. in °C)
26	12	H	2	1	205.3
27	9	F	2	1/2	179.8
28	13	F	3	1	192.5

C. Pharmacological Example10 Example 14 : "apomorphine, tryptamine, norepinephrine (ATN) test in rats"

The antipsychotic activity of the subject compounds is evidenced by the experimental data obtained in the combined apomorphine (APO), tryptamine (TRY) and norepinephrine (NOR) test in rats. Said combined apomorphine, tryptamine and norepinephrine test is described in Arch. Int. Pharmacodyn., 227, 238-253 (1977) and provides an empirical evaluation of the relative specificity with which drugs may effect particular neurotransmitter systems centrally (CNS) as well as peripherally. In particular, the test demonstrates the antagonistic activity of the tested compounds of formula (I) on dopamine (by preventing the symptoms elicited with the dopamine agonist apomorphine), on serotonin (by preventing the central symptoms (convulsions) and peripheral symptoms (hyperaemia) elicited with serotonin or tryptamine), and on

norepinephrine (by preventing or delaying death upon administration of the α -agonist norepinephrine). The favourable pharmacological property of the compounds of formula (I) lies in the absence of a strong α -adrenergic antagonistic activity (column NOR) leading to an improved dissociation between peripheral α -adrenergic antagonistic activity and central serotonin (column TRY convulsions) and central dopamine (column APO) antagonistic activity.

The test is carried out following the procedures described in EP-A-0,196,132 and the experimental data are summarized in Table 5.

10 Table 5

Compound Number	Combined test in rats, ED ₅₀ in mg/kg			
	APO	TRY convulsions	TRY hyperaemia	NOR
1	0.63	0.63	0.04	> 10
2	0.16	0.63	0.01	10
3	0.16	0.16	0.01	> 10
4	0.04	0.04	0.01	2.5
5	0.08	0.31	0.02	> 10
6	0.16	0.16	0.01	10
7	0.63	2.5	0.04	> 10
8	0.63	0.63	0.04	10
10	0.63	0.16	0.0025	10
12	0.31	0.31	0.02	> 10
15	0.16	0.63	0.01	10
16	0.16	0.63	0.02	10
17	0.08	0.16	0.02	5
18	0.08	0.08	0.01	5
19	0.63	2.5	0.63	> 10
20	0.63	0.63	0.16	> 10
24	0.63	0.16	0.01	10
25	0.63	2.5	0.16	> 10
28	0.16	0.16	0.01	> 10
29	0.08	0.31	0.005	5
31	1.25	1.25	0.31	> 10
34	5	> 10	≤ 0.63	≥ 10

D. Composition examples

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I), a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof.

5

Example 15 : Capsules

20 g of the A.I., 6 g sodium lauryl sulfate, 56 g starch, 56 g lactose, 0.8 g colloidal silicon dioxide, and 1.2 g magnesium stearate are vigorously stirred together. The resulting mixture is subsequently filled into 1000 suitable hardened gelatin capsules, each comprising 20 mg of the A.I..

10

Example 16 : Film-coated tabletsPreparation of tablet core

A mixture of 100 g of the A.I., 570 g lactose and 200 g starch is mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture is sieved, dried and sieved again. Then there are added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole is mixed well and compressed into tablets, giving 10,000 tablets, each comprising 10 mg of the active ingredient.

15

Coating

To a solution of 10 g methyl cellulose in 75 ml of denaturated ethanol there is added a solution of 5 g of ethyl cellulose in 150 ml of dichloromethane. Then there are added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene glycol is molten and dissolved in 75 ml of dichloromethane. The latter solution is added to the former and then there are added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated colour suspension and the whole is homogenated. The tablet cores are coated with the thus obtained mixture in a coating apparatus.

20

25

Example 17 : Oral solution

9 Grams of methyl 4-hydroxybenzoate and 1 gram of propyl 4-hydroxybenzoate were dissolved in 4 l of boiling purified water. In 3 l of this solution were dissolved first 10 grams of 2,3-dihydroxybutanedioic acid and thereafter 20 grams of the A.I. The latter solution was combined with the remaining part of the former solution and 12 l 1,2,3-propanetriol and 3 l of sorbitol 70% solution were added thereto. 40 Grams of sodium saccharin were dissolved in 0.5 l of water and 2 ml of raspberry and 2 ml of gooseberry essence were added. The latter solution was combined with the former, water was added q.s. to a volume of 20 l providing an oral solution comprising 5 mg of the

30

35


active ingredient per teaspoonful (5 ml). The resulting solution was filled in suitable containers.

Example 18 : Injectable solution

- 5 1.8 Grams methyl 4-hydroxybenzoate and 0.2 grams propyl 4-hydroxybenzoate were dissolved in about 0.5 l of boiling water for injection. After cooling to about 50°C there were added while stirring 4 grams lactic acid, 0.05 grams propylene glycol and 4 grams of the A.I.. The solution was cooled to room temperature and supplemented with water for injection q.s. ad 1 l, giving a solution comprising 4 mg/ml of A.I.. The solution was
- 10 sterilized by filtration and filled in sterile containers.

Claims

1. A compound of formula



5

a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof, wherein :

the dashed line designates an optional bond;

R¹ and R² are each independently hydrogen, halogen, C₁-6alkyl or C₁-6alkyloxy;


10

R³ and R⁴ are each independently hydrogen, C₁-6alkyl, phenyl or phenyl substituted with one, two or three substituents selected from halo, hydroxy, nitro, cyano, trifluoromethyl, C₁-6alkyl, C₁-6alkyloxy, C₁-6alkylthio, mercapto, amino, mono- and di(C₁-6alkyl)amino, carboxyl, C₁-6alkyloxycarbonyl and C₁-6alkylcarbonyl;

15

Alk is C₁-4alkanediyl;

D is a radical of formula



20

wherein R⁵ is hydrogen or C₁-6alkyl;

R⁶ is hydrogen, C₁-6alkyl, C₁-6alkyloxy, C₁-6alkylthio, amino or mono- or di(C₁-6alkyl)amino; and

R⁷ is hydrogen or C₁-6alkyl; or

25

R⁶ and R⁷ taken together may form a bivalent radical -R⁶-R⁷-, in particular, -R⁶-R⁷- may be

-CH₂-CH₂-CH₂- (a-1);

-CH₂-CH₂-CH₂-CH₂- (a-2);

-CH=CH-CH₂- (a-3);

-CH₂-CH=CH- (a-4) or

30

-CH=CH-CH=CH- (a-5);


wherein one or two hydrogen atoms of said radicals (a-1) to (a-5) each independently may be replaced by C₁-6alkyl, hydroxy, C₁-6alkyloxy or C₁-10alkylcarbonyloxy; or

-R⁶-R⁷- may also be

- 5 -S-CH₂-CH₂- (a-6);
 -S-CH₂-CH₂-CH₂- (a-7);
 -S-CH=CH- (a-8);
 -NH-CH₂-CH₂- (a-9);
 -NH-CH₂-CH₂-CH₂- (a-10);
 10 -NH-CH=CH- (a-11);
 -NH-CH=N- (a-12);
 -S-CH=N- (a-13) or
 -CH=CH-O- (a-14);


15 wherein one or where possible two or three hydrogen atoms in said radicals (a-6) to (a-14) each independently may be replaced by C₁-6alkyl;

or D is a radical of formula




wherein R⁸ is hydrogen or C₁-6alkyl.

2. A compound according to claim 1, wherein D is a radical of formula



-24-



5 wherein R^5 is defined as in claim 1; R^9 and R^{10} each independently are hydrogen or C₁₋₆alkyl; R^{11} is C₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy or C₁₋₁₀alkylcarbonyloxy; and R^{12} is hydrogen, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, amino or mono- or di(C₁₋₆alkyl)amino.

- 10 3. A compound according to claim 1, wherein Alk is C₂₋₄alkanediyl.
4. A compound according to claim 1, wherein R^1 is hydrogen or halogen and R^2 is hydrogen.
- 15 5. A compound according to claim 1, wherein the compound is selected from
 - 3-[2-[4-(5-fluoro-1*H*-indol-1-yl)-1-piperidinyl]ethyl]-6,7,8,9-tetrahydro-2-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;
 - 6-[2-[4-(5-fluoro-1*H*-indol-1-yl)-1-piperidinyl]ethyl]-7-methyl-5*H*-thiazolo-[3,2-*a*]-pyrimidin-5-one;
 - 20 5-[2-[4-(5-fluoro-1*H*-indol-1-yl)-1-piperidinyl]ethyl]-3,6-dimethyl-2-(methylamino)-4(3*H*)-pyrimidinone; and
 - 2,3-dihydro-6-[2-[4-(1*H*-indol-1-yl)-1-piperidinyl]ethyl]-7-methyl-5*H*-thiazolo-[3,2-*a*]pyrimidin-5-one; a pharmaceutically acceptable addition salt or a stereochemically isomeric form thereof.
- 25 6. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective amount of a compound as claimed in any one of the claims 1 to 5.
- 30 7. A process of preparing a pharmaceutical composition as claimed in claim 6, characterized in that a pharmaceutically acceptable carrier is intimately mixed with a therapeutically effective amount of a compound as claimed in any one of claims 1 to 5.

-25-


8. A compound as claimed in any one of claims 1 to 5 for use as a medicine.

9. Use of a compound as claimed in any one of claims 1 to 5 for the manufacture of a medicament for treating psychotic diseases.

5


10. A process of preparing a compound as claimed in claim 1, characterized by

a) *N*-alkylating an intermediate of formula




10 wherein the dashed line, R^1 , R^2 , R^3 and R^4 are defined as in claim 1, with an intermediate of formula D-Alk-W (III), wherein D is defined as in claim 1 and W represents an appropriate reactive leaving group;

b) cyclizing an intermediate of formula



15

wherein D, Alk, R^1 and R^2 are defined as in claim 1 and W is a reactive leaving group; in a reaction inert solvent and in the presence of a reducing agent, thus obtaining a compound of formula



20

wherein D, Alk, R^1 and R^2 are defined as in claim 1;

c) converting compounds of formula (I) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or conversely, converting the acid addition salt form into the free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and, if desired, preparing stereochemically isomeric forms thereof.

25

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 96/00363

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D401/14 C07D487/04 C07D513/04 C07D471/04 C07D495/04
C07D401/04 A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 037 265 (JANSSEN PHARMACEUTICA NV) 7 October 1981 cited in the application see the whole document ---	1-10
A	EP,A,0 070 053 (JANSSEN PHARMACEUTICA NV) 19 January 1983 cited in the application see the whole document ---	1-10
A	EP,A,0 378 255 (JANSSEN PHARMACEUTICA NV) 18 July 1990 cited in the application ---	
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "Z" document member of the same patent family

Date of the actual completion of the international search

31 May 1996

Date of mailing of the international search report

06.06.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Kissler, B

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 96/00363

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 470 039 (LUNDBECK & CO AS H) 5 February 1992 cited in the application see page 5, line 33 - line 37 see claims 1-10	1-10
X	<p>---</p> <p>J. MED. CHEM. (1992), 35(26), 4823-31 CODEN: JMCMAR;ISSN: 0022-2623, 1992, XP002004326 ANDERSEN, KIM ET AL: "Selective, centrally acting serotonin 5-HT₂ antagonists. 2. Substituted 3-(4-fluorophenyl)-1H-indoles" scheme 2, chart 1</p> <p>-----</p>	1-10

INTERNATIONAL SEARCH REPORT
Information on patent family members

International Application No
PCT/EP 96/00363

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0470039		AU-B- 646679	03-03-94
		AU-B- 8141191	06-02-92
		CA-A- 2048027	31-01-92
		DE-D- 69105659	19-01-95
		DE-T- 69105659	27-04-95
		ES-T- 2064974	01-02-95
		HK-A- 51495	13-04-95
		JP-A- 4368367	21-12-92
		NO-B- 178192	30-10-95
		NZ-A- 238956	23-12-93
		US-A- 5393761	28-02-95
